

A rapidly declining perennial sea ice cover in the Arctic

Josefino C. Comiso

Laboratory for Hydrospheric Processes, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

Received 11 June 2002; revised 13 August 2002; accepted 28 August 2002; published 18 October 2002.

[1] The perennial sea ice cover in the Arctic is shown to be declining at -9% per decade using satellite data from 1978 to 2000. A sustained decline at this rate would mean the disappearance of the multiyear ice cover during this century and drastic changes in the Arctic climate system. An apparent increase in the fraction of second year ice in the 1990s is also inferred suggesting an overall thinning of the ice cover. Surface ice temperatures derived from satellite data are negatively correlated with perennial ice area and are shown to be increasing at the rate of 1.2 K per decade. The latter implies longer melt periods and therefore decreasing ice volume in the more recent years. *INDEX TERMS:* 4207 Oceanography: General: Arctic and Antarctic oceanography; 4215 Oceanography: General: Climate and interannual variability (3309); 1635 Global Change: Oceans (4203); 1640 Global Change: Remote sensing. **Citation:** Comiso, J. C., A rapidly declining perennial sea ice cover in the Arctic, *Geophys. Res. Lett.*, 29(20), 1956, doi:10.1029/2002GL015650, 2002.

1. Introduction

[2] The Arctic sea ice cover has been noted as basically impenetrable because of the dominant presence of the perennial ice cover that consists mainly of multiyear ice, the average thickness of which is about 3–4 meters [Wadhams and Comiso, 1992]. The thick multiyear ice floes are the major components of the current Arctic sea ice cover. Their presence during the peak of summer makes a big difference in the ocean-ice-albedo feedback because of their vast extent and high albedo. They survive the summer melt mainly because of a strongly stratified Arctic Ocean that is in part responsible for the scarcity of convection in the region [Aagard and Carmack, 1994]. A study of the perennial ice cover is of immense practical importance because of the potential impact not only on climate but also on the environment and ecology of the system and in light of recent reports of ice retreat [Bjorgo *et al.*, 1997; Parkinson *et al.*, 1999] and ice thinning [Rothrock *et al.*, 1999; Wadhams and Davis, 2000].

[3] In this paper, the state of the perennial sea ice cover is studied using satellite passive microwave data from 1978 through 2000. The multiyear ice cover has been inferred from passive microwave data in winter using a technique that assumes that the signature is spatially stable during this period [Johannessen *et al.*, 1999]. However, previous studies have indicated large regional variations in the passive microwave signature [Grenfell, 1992] causing substantial biases in the derived fraction of multiyear ice within the pack [Kwok *et al.*, 1996]. The key to a more accurate quantification of the multiyear ice cover is through the use

of data during minimum extent since at this time, the seasonal sea ice cover has basically melted and what is left is what we call the perennial ice cover consisting mainly of multiyear ice floes [Comiso, 1990]. These data are also easier to quantify and interpret since no ice classification is needed.

2. Spatial Variability in the Perennial Sea Ice Cover

[4] To provide an overview about interannual variations in the spatial distribution of the perennial ice cover, color-coded ice concentration maps during the summer ice minimum from 1979 to 2000 are shown in Figure 1. The ice concentration maps were derived from satellite passive microwave data using the Bootstrap Algorithm as described in Comiso *et al.* [1997]. Slight adjustments were made to ensure that the data set is temporally consistent using a procedure similar to that used in Parkinson *et al.* [1999]. The day of minimum extent for each year is determined through the use of seven-day running mean data of ice extents. The running mean is used to maximize the chance that what is chosen is the date of real minima and not what might be the result of a temporary compaction due to wind forcing. The dates are mainly in the second or third week of September (Figure 1) and are found to be consistent within a few days with those of ice area minimum.

[5] The images in Figure 1 provide a means to quantitatively identify the relative location and concentration of multiyear ice floes at the end of each ice season. The open water area (blue) around the pack is shown to vary considerably from one year to the next. The circular black areas in the middle are areas not covered by the satellite sensors but are usually highly consolidated. The general location of the perennial sea ice cover changes from one year to another and depends on many factors, the most important of which is the ice drift which has been shown to be strongly influenced by atmospheric circulation [Thorndike and Colony, 1982]. The latter can be in cyclonic mode in which the ice is normally advected to the west causing large open water areas in the east (e.g., Laptev and Kara Seas) and relatively small open areas in the west (e.g., Beaufort Sea and Chukchi Seas) or in anti-cyclonic mode in which the opposite scenario occurs. Examination of the images in Figure 1 also indicates that the open water areas are generally larger in the 1990s than in the 1980s.

[6] To better illustrate the changes from one decade to another, Figure 2a shows the average of the ice concentration minimum maps from 1979 to 1989 while Figure 2b shows the corresponding average from 1990 to 2000. It is apparent that the size of the ice cover in the latter period is smaller than that of the earlier one and that much of the changes occurred around the ice margin. The changes from




Figure 1. Color-coded daily ice concentration maps in the Arctic during ice extent minima for each year from 1979 to 2000. Each map represents the state of the perennial ice cover at the end of the ice season.

one decade to another are better quantified in the difference map (Figure 2c) between the two ice concentration maps. This map shows the magnitude and location of the changes with the negative changes represented by yellows, oranges, purples and reds while positive changes are in grays, greens, and blues. The interdecadal change is surprisingly large and intriguing with the net negative changes in extent and ice area being $5.1 \times 10^5 \text{ km}^2$ (6.3%) and $6.9 \times 10^5 \text{ km}^2$ (11.0%), respectively, during the 22-year period. The biggest change occurred in the western area (Beaufort and Chukchi Seas) while considerable changes are also apparent in the eastern region (Siberian, Laptev and Kara Seas). To get an idea how the ice cover may look like 5 decades from now, the decadal change as reflected in Figure 2c (but shifted towards the north) is applied to the data in Figure 2b and subsequent projections, normalized such that the change is about 10% per decade. The projected perennial ice cover for the decade of the 2050s is shown in Figure 2d. Although the technique is crude and the assumption of a linear decline is likely incorrect, Figure 2d provides a means to assess how the perennial ice cover could look like if the decline persists. It is important to note that the Arctic is governed by complex processes including a positive ice-ocean-atmosphere feedback and decadal as well as interdecadal variability including that associated with the Arctic Oscillation (AO) as described by [Thompson and Wallace, 1998]. A simple regression analysis of AO indices with the perennial ice area indicates that the relationship between these two variables is relatively weak with the

correlation coefficient being only 0.20. The link is likely stronger if continuous ice cover data are analyzed in conjunction with the AO but such study is not within the scope of this paper.

4. Discussion and Conclusions

[13] The area of the Arctic perennial sea ice cover is shown to be declining at a relatively fast rate of $8.9 \pm 2.0\%$ per decade. A decadal change of 10% is also inferred from the difference of 11-year averages of ice minima data. If such a rate of decline persists for a few more decades, the perennial sea ice cover will likely disappear within this century. The decline is unlikely linear because of positive feedback effects between ice, ocean, and the atmosphere. Furthermore, a positive trend in the ice temperature of about 1.2 K per decade is observed leading to earlier onset of melt and delayed onset of freeze up that in turn causes further thinning and retreat of the perennial ice cover.

[14] The implications of such a disappearance of the perennial ice cover are many and can be profound. It would mean a different albedo for the Arctic during the peak of solar insolation in summer and therefore a drastically different ice-ocean-atmosphere feedback. It would mean a much larger influx of solar radiation into the Arctic Ocean thereby changing the characteristics of the mixed layer and the stratification of the ocean. The seasonality and characteristics of the ice cover in the region would be very different. The climate, the productivity, and biota in the region will change tremendously while the region becomes more accessible to human activities.

[15] The Arctic system is however a complex system controlled by many variables and influenced by unpredictable events (e.g., volcanic eruptions). There are also periodic cycles, such as the Arctic Oscillation [Thompson and Wallace, 1998], the North Atlantic Oscillation [Mysak and Venegas, 1998] and the Pacific Decadal Oscillation [Chao et al., 2000] the effects of which need to be considered. The associated decadal and inter-decadal changes in pressure and atmospheric circulation could cause a decadal variability in the ice cover that could lead to a reversal in the current trend. Nevertheless, because of the magnitude in the observed rate of decline and associated feedback effects, a near term recovery is likely needed to avoid an irreversible change in the Arctic ice cover and its environment.