

Ocean primary production and climate: Global decadal changes

Watson W. Gregg

Laboratory for Hydrospheric Processes, NASA/Goddard Space Flight Center, USA

Margarita E. Conkright

Ocean Climate Laboratory, NOAA/National Oceanographic Data Center, USA

Paul Ginoux

NOAA Geophysical Fluid Dynamics Laboratory, USA

John E. O'Reilly

NOAA/National Marine Fisheries Service, USA

Nancy W. Casey

Science Systems and Applications, Inc., USA

Received 7 January 2003; accepted 21 May 2003; published 9 August 2003.

[1] Satellite-in situ blended ocean chlorophyll records indicate that global ocean annual primary production has declined more than 6% since the early 1980's. Nearly 70% of the global decadal decline occurred in the high latitudes. In the northern high latitudes, these reductions in primary production corresponded with increases in sea surface temperature and decreases in atmospheric iron deposition to the oceans. In the Antarctic, the reductions were accompanied by increased wind stress. Three of four low latitude basins exhibited decadal increases in annual primary production. These results indicate that ocean photosynthetic uptake of carbon may be changing as a result of climatic changes and suggest major implications for the global carbon cycle.

INDEX TERMS: 4215 Oceanography: General: Climate and interannual variability (3309); 1615 Global Change: Biogeochemical processes (4805); 1635 Global Change: Oceans (4203); 4806 Oceanography: Biological and Chemical: Carbon cycling. **Citation:** Gregg, W. W., M. E. Conkright, P. Ginoux, J. E. O'Reilly, and N. W. Casey, Ocean primary production and climate: Global decadal changes, *Geophys. Res. Lett.*, 30(15), 1809, doi:10.1029/2003GL016889, 2003.

1. Introduction

[2] Ocean phytoplankton are responsible for approximately half the global biospheric net primary production [Behrenfeld *et al.*, 2001]. Long-term changes in ocean primary production can potentially have important consequences for the global carbon cycle. Global satellite observations of ocean chlorophyll, which are necessary to estimate global ocean primary production, have been available since 1979, but algorithm and processing incompatibilities have precluded time series analysis. Thus an evaluation of decadal changes in global ocean primary production has not previously been possible.

[3] Recently, the historical Coastal Zone Color Scanner (CZCS) record (1979–1986) has been reconstructed [Gregg *et al.*, 2002] to achieve compatibility with the

modern Sea-viewing Wide Field-of-view Sensor (SeaWiFS; 1997-present). In addition, application of blending methodologies [Gregg and Conkright, 2001; Gregg *et al.*, 2002], where in situ data observations are merged with the satellite data, provide improvement of the residual errors of both the CZCS and SeaWiFS. These re-analyzed, blended satellite-in situ chlorophyll data records provide maximum compatibility and permit, for the first time, a quantitative analysis of the changes in global ocean primary production from the early-to-mid 1980's to the present.

2. Methods

[4] Modern methodologies for deriving chlorophyll from space have shown remarkable performance. Eplee *et al.* [2001] showed a mean ratio of SeaWiFS chlorophyll to in situ chlorophyll of 1.0056. Behrenfeld *et al.* [2001] indicated an average difference <6% root mean square. Blending satellite with in situ data can improve the accuracy an additional 4.4% in some basins, depending upon the number of in situ measurements and the size of the error in chlorophyll. Migration of these methodologies to the CZCS data set provides a consistent high quality data record spanning the two decades under investigation here.

[5] SeaWiFS Level-3 Version 4 ocean chlorophyll data were acquired from the Goddard Earth Sciences Distributed Active Archive Center (GES-DAAC) for the period Sep 1997 through Jun 2002. CZCS Level-1 data from Jan 1979 through Jun 1986 were also acquired from the GES-DAAC. The CZCS data set was re-analyzed for compatibility with the SeaWiFS archive [Gregg *et al.*, 2002], including recent algorithm modifications [Stumpf *et al.*, 2003]. Both data sets were blended with in situ data [Gregg and Conkright, 2001] from the NOAA/National Oceanographic Data Center (NODC) and the NASA SeaWiFS Bio-optical Archive and Storage System (SeaBASS).

[6] Global blended chlorophyll seasonal climatologies were used as inputs to the Vertically Generalized Production Model (VGPM) [Behrenfeld and Falkowski, 1997] to compute seasonal ocean primary production. This algorithm is widely used for ocean primary production estimates.




Figure 1. Boundaries for the 12 major oceanographic basins.

Although there are many other algorithms, variability among them associated with chlorophyll, the main driver for decadal changes in primary production, is very small [Campbell *et al.*, 2002]. Seasons were based on the Northern Hemisphere convention, beginning with January through March representing winter. Seasonal analyses were chosen to minimize the effects of data gaps in the CZCS record. The seasonal primary production estimates were integrated to derive annual estimates. The primary production estimates were derived at 1 degree spatial resolution, and coastal regions (bottom depth <200 m) were excluded. Only co-located observations of chlorophyll from CZCS and SeaWiFS were used in the analysis. Also, data poleward of 45° latitude were excluded in boreal spring and summer for the Antarctic, and in boreal autumn and winter in the Atlantic and Pacific to reduce biases associated with poor CZCS sampling in these regions and seasons. Consequently, global annual primary production estimates here are lower than previous estimates [Behrenfeld *et al.*, 2001].

[7] Other inputs to the VGPM are Sea Surface Temperature (SST), daylength, and photosynthetically available radiation (PAR). SST and shortwave radiation data records were obtained from the National Center for Environmental Prediction (NCEP), coincident with the CZCS and SeaWiFS records. The shortwave radiation was converted into PAR (moles quanta $m^{-2} s^{-1}$, over the spectral range 350–700 nm). Decadal changes in SST and PAR had negligible influence on VGPM.

[8] Iron deposition data were taken from a model of mineral dust deposition [Ginoux *et al.*, 2001] and converted to iron assuming a solubility of 2%, and iron content of 5% for a clay fraction, and 1.2% for 3 silt fractions. Only data from 2000–2002 were available to represent the SeaWiFS era, and only data from 1981–1986 were available to represent the CZCS era.

[9] We analyzed SST, surface scalar wind stresses, precipitation, shortwave radiation, sea level pressure, relative humidity, precipitable water, atmospheric iron deposition, aerosol optical depths, and ice cover as potential climatic variables affecting decadal ocean primary production. Only SST, iron deposition, and wind stresses appeared to correspond with the basin-scale changes in primary production and are the only ones shown.

[10] We segregated the oceans into 12 major oceanographic basins (Figure 1) to evaluate basin-scale annual

ocean primary production along with climate change variables. Basin means were area-weighted and only co-located CZCS and SeaWiFS data were used for the comparison. Lower limits for the North Atlantic, North Pacific, and Antarctic were set to 30° N and S to capture the sub-polar frontal zones.

3. Results and Discussion

[11] Global annual ocean primary production has decreased from the CZCS era (1979–1986) to the present by 6.3% ($P < 0.05$; Table 1). SeaWiFS-era (1997–2002) annual primary production was estimated as 42.5 Pg C y^{-1} , compared to 45.3 Pg C y^{-1} for the CZCS era (1979–1986).

[12] The decadal decline in global ocean annual primary production corresponded with an increase in global SST of 0.2°C ($P < 0.05$) over the same time period. Warmer ocean temperatures increase stratification of the surface mixed layer, which inhibits the entrainment of nutrients from below to support ocean primary production [Sarmiento *et al.*, 1998]. In addition, atmospheric iron deposition to the global oceans decreased by 25% over the two observational time segments ($P < 0.05$). Iron is an important micronutrient for phytoplankton and primary production in major parts of the global oceans [Falkowski *et al.*, 1998].

[13] Most of the decrease in global annual primary production occurred in the high latitudes (Figure 2). Combined, the high latitudes losses were 2.0 Pg C y^{-1} , which accounted for 70% of the global decline in photosynthetic carbon uptake. This represented basin-wide decreases of 7, 9, and 10% in the North Atlantic, North Pacific, and Antarctic, respectively, over the two decadal time segments (Table 1). Concurrently, the North Atlantic and North Pacific experienced major increases in SST: 0.7 and 0.4°C respectively (Figure 2). These reductions in primary production and accompanying increases in SST may be related via increased stratification and reduced nutrient supply, as described above. Although increased temperature and shallower mixed layers can lead to increased chlorophyll in early spring in these basins, more severe nutrient exhaustion reduces concentrations in late spring and summer [Gregg and Conkright, 2002] and results in reduced annual primary production. Similar long-term reductions in primary production (along with decreasing availability of nutrients) have been observed in the North Pacific by other investigators [Ono *et al.*, 2002].

Table 1. Percent change in ocean primary production (SeaWiFS-CZCS) by basin and surface area of basins ($10^7 km^{-2}$) where data from both SeaWiFS and CZCS were sampled

Basin	Area	Change
N. Atlantic	1.83	−6.7%
N. Pacific	2.32	−9.3%
N. Central Atlantic	1.53	−7.0%
N. Central Pacific	3.22	−5.8%
N. Indian	0.46	13.6%
Eq. Atlantic	1.15	6.9%
Eq. Pacific	3.72	−3.0%
Eq. Indian	1.37	8.8%
S. Atlantic	1.20	−3.8%
S. Pacific	2.69	−14.0%
S. Indian	1.77	−4.2%
Antarctic	8.28	−10.4%
Global	29.73	−6.3%